

# Differential Sandwich theorems of $p$ -valent functions defined by Noor integral operator

Ping He\* , Defei Zhang

Department of Mathematics  
Honghe University  
Mengzi, 661100, Yunnan, China  
e-mail: hepingky@163.com

## Abstract

*In this paper, we obtain sandwich results involving Hadamard product for certain  $p$ -valent functions associated with Noor integral operator in the open unit disk.*

**Keywords:** *Sandwich theorems; Hadamard product; Operator.*

**2000 Mathematical Subject Classification:** 47A06.

## 1 Introduction

Let  $\Sigma_p$  denote the class of functions  $f(z)$  of the form

$$f(z) = z^p + \sum_{k=p+1}^{\infty} a_k z^k \quad (p \in N = 1, 2, 3, \dots), \quad (1)$$

which are analytic in the punctured open unit disk  $\mathbb{U} = \{z : |z| < 1\}$ .

For functions  $f \in \Sigma_p$  given by (1) and  $g \in \Sigma_p$  given by

$$g(z) = z^p + \sum_{k=p+1}^{\infty} b_k z^k.$$

We define the Hadamard product (or convolution) of  $f$  and  $g$  by

$$(f * g)(z) = z^p + \sum_{k=p+1}^{\infty} a_k b_k z^k. \tag{2}$$

Let  $f(z)$  and  $g(z)$  be analytic in  $\mathbb{U}$ . We say that the function  $g(z)$  is subordinate to  $f(z)$ , if there exists a function  $w(z)$  analytic in  $\mathbb{U}$ , with  $w(0) = 0$  and  $|w(z)| < 1$ , and such that  $g(z) = f(w(z))$ . In such a case, we write  $g(z) \prec f(z)$ . If the function  $f$  is univalent in  $\mathbb{U}$ , then  $g(z) \prec f(z)$  if and only if  $g(0) = f(0)$  and  $g(\mathbb{U}) \subset f(\mathbb{U})$ .

Let  $H(\mathbb{U})$  denote the class of analytic functions in  $\mathbb{U}$  and let  $H(a, n)$  denote the subclass of functions  $f \in H(\mathbb{U})$  of the form:

$$f(z) = a + a_n z^n + a_{n+1} z^{n+1} + \dots (n = 1, 2, 3, \dots).$$

Denote by  $Q$ , the set of all functions  $f(z)$  that are analytic and injective on  $\overline{U} \setminus E(f)$ , where  $E(f) = \{\xi \in \partial U : \lim_{z \rightarrow \xi} f(z) = \infty\}$ , and such that  $f'(\xi) \neq 0$  for  $\xi \in \partial U \setminus E(f)$ .

Let  $\psi : \mathbb{C}^3 \times \overline{U} \rightarrow \mathbb{C}$ , let  $h(z)$  be univalent in  $\mathbb{U}$  and  $q(z) \in Q$ . Miller and Mocanu[1] considered the problem of determining conditions on admissible function  $\psi$  such that

$$\psi(p(z), zp'(z), z^2 p''(z); z) \prec h(z) \tag{3}$$

implies  $p(z) \prec q(z)$ , for all functions  $p(z) \in H(a, n)$  that satisfy the differential subordination (3). Moreover, they found conditions so that  $q(z)$  is the smallest function with this property, called the best dominant of the subordination (3).

Let  $\varphi : \mathbb{C}^3 \times \overline{U} \rightarrow \mathbb{C}$ , let  $h(z) \in H$  and  $q(z) \in H(a, n)$ . Recently Miller and Mocanu[2] studied the dual problem and determined conditions on  $\varphi$  such that

$$h(z) \prec \varphi(p(z), zp'(z), z^2 p''(z); z) \tag{4}$$

implies  $q(z) \prec p(z)$ , for all functions  $p(z) \in Q$  that satisfy the above superordination. They also found conditions so that the function  $q(z)$  is the largest function with this property, called the best subordinant of the superordination (4).

Liu and Noor [3] introduced an integral operator  $\mathcal{N}_{n,p} f(z) : \Sigma_p \rightarrow \Sigma_p$  as follows:

Let  $f_{n,p}(z) = \frac{z^p}{(1-z)^{n+p}}$  ( $n > -p$ ), and let  $f_{n,p}^{(+)}(z)$  be defined such that

$$f_{n,p}(z) * f_{n,p}^{(+)}(z) = \frac{z^p}{(1-z)^{1+p}}, \tag{5}$$

then

$$\mathcal{N}_{n,p} f(z) = f_{n,p}^{(+)}(z) * f_{n,p}(z) = \left(\frac{z^p}{(1-z)^{n+p}}\right)^{(+)} * f_{n,p}(z). \tag{6}$$

If  $f(z)$  is given by (1), then from (5) and (6), we deduce that

$$\mathcal{N}_{n,p}f(z) = z^p + \sum_{k=p+1}^{\infty} \frac{(p+1)(p+2)\cdots k}{(n+p)(n+p+1)\cdots(n+k-1)} a_k z^k. \quad (7)$$

It follows from (7) that

$$z(\mathcal{N}_{n+1,p}f(z))' = (n+p)\mathcal{N}_{n,p}f(z) - n\mathcal{N}_{n+1,p}f(z). \quad (8)$$

We also note that  $\mathcal{N}_{0,p}f(z) = \frac{zf'(z)}{p}$  and  $\mathcal{N}_{1,p}f(z) = f(z)$ . The operator  $\mathcal{N}_{n,p}f(z)$  defined by (6) is called as the Noor integral operator of  $(n+p-1)$ th order of  $f(z)$ [1]. For  $p=1$ , the operator  $\mathcal{N}_{n,1}f(z)$  was introduced by Noor[4] and Noor[5]. Several classes of analytic functions, defined by using the operator  $\mathcal{N}_{n,1}f(z)$ , have been studied by many authors[6,7,8].

In this paper, we will derive several subordination results, superordination results and sandwich results involving the operator  $\mathcal{N}_{n,p}f(z)$  and some of its special operators.

In order to prove our main results, we need the following lemmas.

**Lemma 1** (see [9]). Let  $q(z)$  be univalent in  $\mathbb{U}$ ,  $\gamma \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}$  and suppose that

$$\operatorname{Re}\left\{1 + \frac{zq''(z)}{q'(z)}\right\} > \max\left\{0, -\operatorname{Re}\frac{1}{\gamma}\right\}. \quad (9)$$

If  $p(z)$  is analytic in  $\mathbb{U}$ , with  $p(0) = q(0)$  and

$$p(z) + \gamma zp'(z) \prec q(z) + \gamma zq'(z), \quad (10)$$

then  $p(z) \prec q(z)$ , and  $q(z)$  is the best dominant.

**Lemma 2** (see [10]). Let  $q(z)$  be convex in  $\mathbb{U}$ ,  $q(0) = a$  and  $\gamma \in \mathbb{C}$ ,  $\operatorname{Re}\gamma > 0$ . If  $p \in H(a, 1)$  and  $p(z) + \gamma zp'(z)$  is univalent in  $\mathbb{U}$ , and

$$q(z) + \gamma zq'(z) \prec p(z) + \gamma zp'(z), \quad (11)$$

then  $q(z) \prec p(z)$  and  $q(z)$  is the best subdominant.

## 2 Main Results

We shall assume in the remainder of this paper that  $p, n \in \mathbb{N}$  and  $z \in \mathbb{U}$ .

**Theorem 1.** Let  $q(z)$  be univalent in  $\mathbb{U}$  with  $q(0) = 1$ ,  $\alpha \in \mathbb{C}^*$ , and suppose that

$$\operatorname{Re}\left\{1 + \frac{zq''(z)}{q'(z)}\right\} > \max\left\{0, -\operatorname{Re}\frac{n+p}{\alpha}\right\}. \quad (12)$$

If  $f(z) \in \Sigma_p$  satisfies the subordination

$$\mathcal{R}(\alpha, n, p) \prec q(z) + \frac{\alpha}{n+p} zq'(z), \tag{13}$$

where  $\mathcal{R}(\alpha, n, p)$  is given by

$$\mathcal{R}(\alpha, n, p) = (1 - \alpha) \frac{\mathcal{N}_{n+1,p}f(z)}{z^p} + \alpha \frac{\mathcal{N}_{n,p}f(z)}{z^p}, \tag{14}$$

then

$$\frac{\mathcal{N}_{n+1,p}f(z)}{z^p} \prec q(z).$$

and  $q(z)$  is the best dominant.

**Proof .** Let

$$p(z) = \frac{\mathcal{N}_{n+1,p}f(z)}{z^p}, \tag{15}$$

differentiating (15) with respect to  $z$  and using the identity (8) in the resulting equation, we have

$$\frac{zp'(z)}{p(z)} = (n+p) \left\{ \frac{\mathcal{N}_{n,p}f(z)}{\mathcal{N}_{n+1,p}f(z)} - 1 \right\},$$

that is,

$$\frac{1}{(n+p)} zp'(z) = \frac{\mathcal{N}_{n,p}f(z)}{z^p} - \frac{\mathcal{N}_{n+1,p}f(z)}{z^p}.$$

Therefore, we have

$$\mathcal{R}(\alpha, n, p) = (1 - \alpha) \frac{\mathcal{N}_{n+1,p}f(z)}{z^p} + \alpha \frac{\mathcal{N}_{n,p}f(z)}{z^p} = p(z) + \frac{\alpha}{(n+p)} zp'(z).$$

By (13), we obtain

$$p(z) + \frac{\alpha}{(n+p)} zp'(z) \prec q(z) + \frac{\alpha}{n+p} zq'(z).$$

By Lemma 1,  $\frac{\mathcal{N}_{n+1,p}f(z)}{z^p} \prec q(z)$ , and the proof of Theorem 1 is completed.

Taking the convex function  $q(z) = \frac{1+Az}{1+Bz}$  in Theorem 1, we have the following corollary.

**Corollary 1.** Let  $A, B, \alpha \in \mathbb{C}$ ,  $A \neq B$ ,  $|B| < 1$ ,  $Re\alpha > 0$ . If  $f(z) \in \Sigma_p$  satisfies the subordination

$$\mathcal{R}(\alpha, n, p) \prec \frac{1+Az}{1+Bz} + \frac{\alpha}{n+p} \frac{(A-B)z}{(1+Bz)^2},$$

where  $\mathcal{R}(\alpha, n, p)$  is given by (14), then

$$\frac{\mathcal{N}_{n+1,p}f(z)}{z^p} \prec \frac{1+Az}{1+Bz},$$

and the function  $\frac{1+Az}{1+Bz}$  is the best dominant.

Taking  $n = 0$  in Theorem 1, we obtain the following result.

**Corollary 2.** Let  $q(z)$  be univalent in  $\mathbb{U}$  with  $q(0) = 1$ ,  $\alpha \in \mathbb{C}^*$ , and suppose that (12) holds. If  $f(z) \in \Sigma_p$  satisfies the subordination

$$\mathcal{R}(\alpha, p) \prec q(z) + \frac{\alpha}{p} zq'(z), \quad (16)$$

where  $\mathcal{R}(\alpha, p)$  is given by

$$\mathcal{R}(\alpha, p) = (1 - \alpha) \frac{f(z)}{z^p} + \frac{\alpha}{p} \frac{zf'(z)}{z^p}, \quad (17)$$

then

$$\frac{f(z)}{z^p} \prec q(z).$$

Taking  $p = 1$  in Theorem 1, we have the following result.

**Corollary 3.** Let  $q(z)$  be univalent in  $\mathbb{U}$  with  $q(0) = 1$ ,  $\alpha \in \mathbb{C}^*$ , and suppose that (12) holds. If  $f(z) \in \Sigma_p$  satisfies the subordination

$$\mathcal{R}(\alpha, n) \prec q(z) + \frac{\alpha}{n+1} zq'(z), \quad (18)$$

where  $\mathcal{R}(\alpha, n)$  is given by

$$\mathcal{R}(\alpha, n) = (1 - \alpha) \frac{\mathcal{N}_{n+1}f(z)}{z} + \alpha \frac{\mathcal{N}_n f(z)}{z}, \quad (19)$$

then

$$\frac{\mathcal{N}_{n+1}f(z)}{z} \prec q(z).$$

**Theorem 2.** Let  $q(z)$  be convex in  $\mathbb{U}$ ,  $q(0) = 1$  and  $\alpha \in \mathbb{C}$ ,  $Re\alpha > 0$ . If  $f(z) \in \Sigma_p$  such that  $\frac{\mathcal{N}_{n+1,p}f(z)}{z^p} \in H(q(0), 1) \cap Q$ , and  $\mathcal{R}(\alpha, n, p)$  is univalent in  $\mathbb{U}$  and satisfies the superordination

$$q(z) + \frac{\alpha}{n+p} zq'(z) \prec \mathcal{R}(\alpha, n, p), \quad (20)$$

where  $\mathcal{R}(\alpha, n, p)$  is given by (14), then

$$q(z) \prec \frac{\mathcal{N}_{n+1,p}f(z)}{z^p},$$

and  $q(z)$  is the best subordinator.

**Proof .** Let  $p(z)$  be given by (15) and proceeding as in the proof of Theorem 1, the subordination (20) becomes

$$q(z) + \frac{\alpha}{n+p} zq'(z) \prec p(z) + \frac{\alpha}{(n+p)} zp'(z).$$

The proof follows by an application of Lemma 2.

Taking  $n = 0$  in Theorem 2, we obtain the following result.

**Corollary 4.** Let  $q(z)$  be convex in  $\mathbb{U}$ ,  $q(0) = 1$  and  $\alpha \in \mathbb{C}$ ,  $Re\alpha > 0$ . If  $f(z) \in \Sigma_p$  such that  $\frac{f(z)}{z^p} \in H(q(0), 1) \cap Q$ , and  $\mathcal{R}(\alpha, p)$  is univalent in  $\mathbb{U}$  and satisfies the superordination

$$q(z) + \frac{\alpha}{p} zq'(z) \prec \mathcal{R}(\alpha, p), \quad (21)$$

where  $\mathcal{R}(\alpha, p)$  is given by(17), then

$$q(z) \prec \frac{f(z)}{z^p}.$$

Taking  $p = 1$  in Theorem 2, we have the following result.

**Corollary 5.** Let  $q(z)$  be convex in  $\mathbb{U}$ ,  $q(0) = 1$  and  $\alpha \in \mathbb{C}$ ,  $Re\alpha > 0$ . If  $f(z) \in \Sigma_p$  such that  $\frac{\mathcal{N}_{n+1}f(z)}{z} \in H(q(0), 1) \cap Q$ , and  $\mathcal{R}(\alpha, n)$  is univalent in  $\mathbb{U}$  and satisfies the superordination

$$\mathcal{R}(\alpha, n) \prec q(z) + \frac{\alpha}{n+1} zq'(z), \quad (22)$$

where  $\mathcal{R}(\alpha, n)$  is given by (19), then

$$q(z) \prec \frac{\mathcal{N}_{n+1}f(z)}{z}.$$

Combining Theorems 1 and 2, we have the following sandwich theorem.

**Theorem 3.** Let  $q_1$  and  $q_2(z)$  be convex in  $\mathbb{U}$ ,  $q_1(0) = q_2(0) = 1$  and  $q_2(z)$  satisfies (12), and  $\alpha \in \mathbb{C}$ ,  $Re\alpha > 0$ . If  $f(z) \in \Sigma_p$  such that  $\frac{\mathcal{N}_{n+1,p}f(z)}{z} \in H(q_1(0), 1) \cap Q$ , and  $\mathcal{R}(\alpha, n, p)$  is univalent in  $\mathbb{U}$  and satisfies

$$q_1(z) + \frac{\alpha}{n+p} zq_1'(z) \prec \mathcal{R}(\alpha, n, p) \prec q_2(z) + \frac{\alpha}{n+p} zq_2'(z), \quad (23)$$

where  $\mathcal{R}(\alpha, n, p)$  is given by (14), then

$$q_1(z) \prec \frac{\mathcal{N}_{n+1,p}f(z)}{z^p} \prec q_2(z).$$

and  $q_1(z)$ ,  $q_2(z)$  are the best subordinant and the best dominant, respectively.

Remark . Combining Corollaries 2, 4 and Corollaries 3, 5, we obtain the corresponding sandwich results for the operators  $\mathcal{N}_p$  and  $\mathcal{N}_{n+1}$

### 3 Open Problem

If  $f(z)$  is meromorphically multivalent functions, a new operator can be defined. New results which is about differential Sandwich theorems of the new operator can be obtained.

## 4 Acknowledgements

This work was supported by the Natural Science Foundation of Yunnan(2013FZ116) ,and the Scientific Research Foundation from Yunnan Province Education Committee (2010Y167)and (2011C120), and the Quality Engineering Construction of Honghe University (ZDKC 1111) .

## References

- [1] S.S. Miller, P.T. Mocanu, Differential Subordination: Theory and Applications, in: Series on Monographs and Textbooks in Pure and Applied Mathematics, vol. 225, Marcel Dekker Inc, New York, Basel, 2000.
- [2] S.S. Miller, P.T. Mocanu, Subordinates of differential subordinations, *Complex Var.* 48 (10) (2003) 815-826.
- [3] J.-L. Liu, K.I. Noor, Some properties of Noor integral operator, *J. Natur. Geom.* 21 (2002) 81-90.
- [4] K.I. Noor, On new classes of integral operators, *J. Natur. Geom.* 16 (1999) 71-80.
- [5] K.I. Noor, M.A. Noor, On integral operators, *J. Math. Anal. Appl.* 238 (1999) 341-352.
- [6] N.E. Cho, The Noor integral operator and strongly close-to-convex functions, *J. Math. Anal. Appl.* 283 (2003) 202-212.
- [7] J.-L. Liu, The Noor integral and strongly starlike functions, *J. Math. Anal. Appl.* 261 (2001) 441-447.
- [8] K.I. Noor, M.A. Noor, On certain classes of analytic functions defined by Noor integral operator, *J. Math. Anal. Appl.* 281 (2003) 244-252.
- [9] L. Cotirlă, A differential sandwich theorem for analytic functions defined by the integral operator, *Stud. Univ. Babeş -BolyaiMath.* 54 (2) (2009) 13-21.
- [10] T. Bulboacă, Classes of first order differential subordinations, *Demonstratio Math.* 35 (2) (2002) 287-292.